



SCANNED TO PHARMACY

STAT MEDICATION

Peripherally Inserted Central Catheter (PICC) Orders

1. IV Therapy PICC nurse to Evaluate for and Insert PICC line as indicated.
2. IV Therapy PICC nurse unable to insert PICC: consult to Interventional Radiology for Venous Access.
3. Lidocaine 1% 1–3 ml intradermal per procedure.
4. If PICC inserted by IV Therapy, order:
 - _____ STAT Chest X-ray for PICC Tip placement, call report to IV Therapy floor 1, 2 or 3 via Vocera.
 - _____ STAT portable Chest X-ray for PICC Tip placement, call report to IV Therapy floor 1, 2 or 3 via Vocera.
5. Implement PICC Protocol for flushing, access and care of PICC line.
6. K-Pad to PICC Right/Left arm continuous for 24 hours, then PRN.
7. Alteplase (Tissue Plasminogen Activator) 2mg/2ml to each catheter lumen per catheter clearance protocol after IV Nurse Specialist evaluation. May repeat X 1. May be administered by IVT RN or RN completing TPA competency.

Authorization for therapeutic substitution is given unless checked here

Date _____ Time _____ Physician _____



PICC Orders – Adult

Patient Identification

11-21-07
R006 / 50-7715-0572
63

File Specification Information

Document Size	Gripper / Lock-Up	Stub	Imprint
8.5 x 11	ALL	N	NA
Punch	Numbering	Binding	Registering Plys
5 T	N	N	N
Screens	Backer(s) and Percentage		
N	N		
Color(s)	BLK		
File Information / Master Page			
Pyxis Format			
Other			